

# ON THOMPSON'S $p$ -COMPLEMENT THEOREMS FOR SATURATED FUSION SYSTEMS

JON GONZÁLEZ-SÁNCHEZ, ALBERT RUIZ, AND ANTONIO VIRUEL

ABSTRACT. In this short note we prove that a saturated fusion system admitting some special type of automorphism is nilpotent. This generalizes classical results by J.G. Thompson.

## 1. INTRODUCTION

In his PhD Thesis, John G. Thompson proved the long standing conjecture that the Frobenius kernel of a Frobenius group is nilpotent [Tho59]. The Frobenius kernel always admits a fixed-point-free automorphism of prime order which turned out to be a sufficient condition for the nilpotency. In fact both results, the nilpotency of the Frobenius kernel and the nilpotency of a finite group admitting an fixed-point-free automorphism of prime order are equivalent. In order to prove this result, Thompson introduced his famous  $p$ -nilpotency criterion for an odd prime  $p$ , a group  $G$  is  $p$ -nilpotent if and only if the normalizer of the  $J$  subgroup of a Sylow  $p$ -subgroup and the centralizer of the center of a Sylow  $p$ -subgroup are  $p$ -nilpotent (see [Tho64]). In fact, Thompson proved that a group  $G$  admitting automorphisms that leave invariant some special subgroups is  $p$ -nilpotent. The following translates [Tho60, Theorem A] for saturated fusion systems, and includes an extra hypothesis to cover the  $p = 2$  case:

**Theorem 1.1.** *Let  $\mathcal{F}$  be a saturated fusion system over a  $p$  group  $S$  such that either  $p$  is odd, or  $p = 2$  and  $\mathcal{F}$  is  $\Sigma_4$ -free. Let  $\mathcal{U}$  be a group of automorphisms of  $(S, \mathcal{F})$ . Suppose for every  $\mathcal{U}$ -invariant normal subgroup  $Q \trianglelefteq S$ ,  $\text{Aut}_{\mathcal{F}}(Q)$  is a  $p$ -group. Then  $\mathcal{F} = \mathcal{F}_S(S)$ .*

As a consequence of [Tho60, Theorem A] Thompson proved that a group  $G$  admitting a fixed-point-free automorphism of prime order is  $p$ -nilpotent for all primes, and therefore the group is nilpotent [Tho59, Theorem 1].

Given a finite group  $G$  and  $\phi$  a prime order automorphism without fix points, then  $\phi$  fixes a Sylow  $p$ -subgroup  $S$  of  $G$ . For a fixed subgroup  $H$  one has that  $N_G(H)$  and  $C_G(H)$  are also fixed. Therefore,  $\phi$  acts on the quotient group  $N_G(H)/C_G(H)$  and the action on this quotient is without fix points (see Lemma 2.9(b)). So one

---

2010 *Mathematics Subject Classification.* 20D20, 55R35.

First author is partially supported by MINECO MTM2011-28229-C02-01. Second author is partially supported by FEDER-MICINN grant MTM2010-20692. Third author is partially supported by FEDER-MICINN grant MTM2010-18089, and Junta de Andalucía grants FQM-213 and P07-FQM-2863. Second and third authors are partially supported by Generalitat de Catalunya grant 2009SGR-1092.

can translate the concept of a fixed-point-free automorphism of prime order from the category of finite groups to the category of saturated fusion systems (see Definition 2.10 and Proposition 2.11). It turned out that the concept of fixed-point-free automorphism in the category of saturated fusion systems is more general since there exist finite groups admitting automorphism of prime order with fix points such that the induced automorphism on the fusion category is fixed-point-free (see Example 3.4). Nevertheless,  $p$ -nilpotency holds under this mild assumption, as we prove this generalization of [Tho59, Theorem 1]:

**Theorem 1.2.** *Let  $\mathcal{F}$  be a saturated fusion system over a  $p$  group  $S$  such that either  $p$  is odd, or  $p = 2$  and  $\mathcal{F}$  is  $\Sigma_4$ -free. Then, if  $(S, \mathcal{F})$  admits a prime order fixed-point-free automorphism,  $\mathcal{F} = \mathcal{F}_S(S)$ .*

These results contribute to the list of nilpotency criteria for saturated fusion systems that generalize classical criteria for finite groups, and fits within the framework of previous work by Kessar-Linckelmann [KL08], Díaz-Glesser-Park-Stancu [DGPS11] and Cantarero-Scherer-Viruel [CSV13].

The note is organized as follows. In Section 2 we recall the main properties of saturated fusion system and we introduce the concept of a fixed-point-free automorphism of a saturated fusion system. In Section 3 we provide a unified proof of Theorems 1.1 and 1.2.

## 2. BACKGROUND ON SATURATED FUSION SYSTEMS AND FINITE GROUPS

In this section we review the concept of a saturated fusion system over a  $p$ -group  $S$  as defined in [BLO03], and define fixed-point-free automorphism of a saturated fusion system.

**Definition 2.1.** A fusion system  $\mathcal{F}$  over a finite  $p$ -group  $S$  is a category whose objects are the subgroups of  $S$ , and whose morphisms sets  $\text{Hom}_{\mathcal{F}}(P, Q)$  satisfy the following two conditions:

- (a)  $\text{Hom}_S(P, Q) \subseteq \text{Hom}_{\mathcal{F}}(P, Q) \subseteq \text{Inj}(P, Q)$  for all  $P$  and  $Q$  subgroups of  $S$ .
- (b) Every morphism in  $\mathcal{F}$  factors as an isomorphism in  $\mathcal{F}$  followed by an inclusion.

We say that two subgroups  $P, Q \leq S$  are  $\mathcal{F}$ -conjugate if there is an isomorphism between them in  $\mathcal{F}$ . As all the morphisms are injective by condition (b), we denote by  $\text{Aut}_{\mathcal{F}}(P)$  the group  $\text{Hom}_{\mathcal{F}}(P, P)$ . We denote by  $\text{Out}_{\mathcal{F}}(P)$  the quotient group  $\text{Aut}_{\mathcal{F}}(P) / \text{Aut}_P(P)$ .

The fusion systems that we consider are saturated, so we need the following definitions:

**Definition 2.2.** Let  $\mathcal{F}$  be a fusion system over a  $p$ -group  $S$ .

- A subgroup  $P \leq S$  is *fully centralized in  $\mathcal{F}$*  if  $|C_S(P)| \geq |C_S(P')|$  for all  $P'$  which is  $\mathcal{F}$ -conjugate to  $P$ .
- A subgroup  $P \leq S$  is *fully normalized in  $\mathcal{F}$*  if  $|N_S(P)| \geq |N_S(P')|$  for all  $P'$  which is  $\mathcal{F}$ -conjugate to  $P$ .
- $\mathcal{F}$  is a *saturated fusion system* if the following two conditions hold:

- (I) Every fully normalized in  $\mathcal{F}$  subgroup  $P \leq S$  is fully centralized in  $\mathcal{F}$  and  $\text{Aut}_S(P) \in \text{Syl}_p(\text{Aut}_{\mathcal{F}}(P))$ .
- (II) If  $P \leq S$  and  $\varphi \in \text{Hom}_{\mathcal{F}}(P, S)$  are such that  $\varphi P$  is fully centralized, and if we set

$$N_{\varphi} = \{g \in N_S(P) \mid \varphi c_g \varphi^{-1} \in \text{Aut}_S(\varphi P)\},$$

then there is  $\bar{\varphi} \in \text{Hom}_{\mathcal{F}}(N_{\varphi}, S)$  such that  $\bar{\varphi}|_P = \varphi$ .

As expected, every finite group  $G$  gives rise to a saturated fusion system over  $S$ , a Sylow  $p$ -subgroup of  $G$ , denoted by  $(S, \mathcal{F}_G(S))$  [BLO03, Proposition 1.3]. But there exist saturated fusion systems that are not the fusion system of any finite group, e.g. [BLO03, Section 9] or [RV04].

Let  $\mathcal{F}$  be a fusion system over a  $p$ -group  $S$ , and  $Q$  a subgroup of  $S$ . We can take the *normalizer of  $Q$  in  $\mathcal{F}$*  as the fusion system over the normalizer of  $Q$  in  $S$ ,  $N_S(Q)$ , with morphisms:

$$\text{Hom}_{N_{\mathcal{F}}(Q)}(P, P') = \{\varphi \in \text{Hom}_{\mathcal{F}}(P, P') \mid \exists \psi \in \text{Hom}_{\mathcal{F}}(PQ, P'Q), \psi|_P = \varphi\}.$$

Although,  $(N_S(Q), N_{\mathcal{F}}(Q))$  is not always a saturated fusion system, it is so when  $Q$  is fully normalized [BLO03, Proposition A.6]:

**Proposition 2.3.** *Let  $\mathcal{F}$  be a saturated fusion system over a  $p$ -group  $S$ . If  $Q \leq S$  is fully normalized in  $\mathcal{F}$ , then  $(N_S(Q), N_{\mathcal{F}}(Q))$  is a saturated fusion system.*

We also need results concerning the quotients of saturated fusion systems. Recall that if  $(S, \mathcal{F})$  is a saturated fusion system, we say that  $Q \leq S$  is a *weakly  $\mathcal{F}$ -closed* subgroup if  $Q$  is not  $\mathcal{F}$ -conjugate to any other subgroup of  $S$ .

In [Oli04, Lemma 2.6] we can find the following result:

**Lemma 2.4.** *Let  $(S, \mathcal{F})$  be a saturated fusion system, and let  $Q \triangleleft S$  be a weakly  $\mathcal{F}$ -closed subgroup. Let  $\mathcal{F}/Q$  be the fusion system over  $S/Q$  defined by setting:*

$$\text{Hom}_{\mathcal{F}/Q}(P/Q, P'/Q) = \{\varphi/Q \mid \varphi \in \text{Hom}_{\mathcal{F}}(P, P')\}$$

for all  $P, P' \leq S$  which contains  $Q$ . Then  $\mathcal{F}/Q$  is saturated.

Some classical results for finite groups can be generalized to saturated fusion systems, as for example, Alperin's Fusion Theorem for saturated fusion systems [BLO03, Theorem A.10]:

**Definition 2.5.** Let  $\mathcal{F}$  be any fusion system over a  $p$ -group  $S$ . A subgroup  $P \leq S$  is:

- $\mathcal{F}$ -centric if  $P$  and all its  $\mathcal{F}$ -conjugates contain their  $S$ -centralizers.
- $\mathcal{F}$ -radical if  $\text{Out}_{\mathcal{F}}(P)$  is  $p$ -reduced, that is, if  $\text{Out}_{\mathcal{F}}(P)$  has no nontrivial normal  $p$ -subgroups.

**Theorem 2.6.** *Let  $\mathcal{F}$  be a saturated fusion system over  $S$ . Then for each morphism  $\psi \in \text{Aut}_{\mathcal{F}}(P, P')$ , there exists a sequence of subgroups of  $S$*

$$P = P_0, P_1, \dots, P_k = P' \quad \text{and} \quad Q_1, Q_2, \dots, Q_k,$$

and morphisms  $\psi_i \in \text{Aut}_{\mathcal{F}}(Q_i)$ , such that

- $Q_i$  is fully normalized in  $\mathcal{F}$ ,  $\mathcal{F}$ -radical and  $\mathcal{F}$ -centric for each  $i$ ;
- $P_{i-1}, P_i \leq Q_i$  and  $\psi_i(P_{i-1}) = P_i$  for each  $i$ ; and
- $\psi = \psi_k \circ \psi_{k-1} \circ \cdots \circ \psi_1$ .

Here we will recall some definitions and results concerning groups [Gor68, Chapter 2] and group automorphisms [Gor68, Chapter 10]:

A finite group  $G$  is *nilpotent* if the lower central series of  $G$ , defined as

$$\gamma_1(G) = G, \gamma_i(G) = [\gamma_{i-1}(G), G] \text{ for } i \geq 2$$

satisfies that there exists  $m$  such that  $\gamma_m = \{1\}$ .

As examples we have that every finite  $p$ -group is nilpotent [Gor68, Theorem 3.3(iii)], and more generally:

**Theorem 2.7.** *A finite group  $G$  is nilpotent if and only if it is the direct product of its Sylow  $p$ -subgroups.*

So the fusion on  $S$ , a Sylow  $p$ -subgroup of a nilpotent group  $G$ , satisfies  $\mathcal{F}_S(G) = \mathcal{F}_S(S)$ .

**Definition 2.8.** An automorphism  $\varphi$  of a group  $G$  is said to be *fixed-point-free* if it leaves only the identity element of  $G$  fixed.

The following result shows that fixed-point-free are compatible with the  $p$ -local structure [Gor68, Theorem 10.1.2 and Lemma 10.1.3]:

**Theorem 2.9.** *Let  $G$  be a finite group and  $p$  a prime dividing the order of  $G$ . If  $\varphi: G \rightarrow G$  is a fixed-point-free morphism then:*

- There exists a unique  $\varphi$ -invariant Sylow  $p$ -subgroup  $S$  of  $G$ , and it contains every  $\varphi$ -invariant  $p$ -subgroup of  $G$ .*
- If  $H$  is a  $\varphi$ -invariant normal subgroup of  $G$ , then  $\varphi$  induces a fixed-point-free automorphism of  $G/H$ .*

Let  $\text{Aut}(\mathcal{F})$  denote the group of automorphisms of  $\mathcal{F}$ :

$$\text{Aut}(\mathcal{F}) = \{\varphi \in \text{Aut}(S) \mid \text{if } \alpha \in \text{Hom}_{\mathcal{F}}(P, Q)$$

$$\text{then } \varphi|_Q \circ \alpha \circ (\varphi|_P)^{-1} \in \text{Hom}_{\mathcal{F}}(\varphi(P), \varphi(Q))\}.$$

We are now ready to give a definition of what a fixed-point-free automorphism of a saturated fusion system is.

**Definition 2.10.** Let  $(S, \mathcal{F})$  be a saturated fusion system. Then  $\varphi \in \text{Aut}(\mathcal{F})$  is a *fixed-point-free* automorphism if the following hold

- $\varphi: S \rightarrow S$  is fixed-point-free and
- $\varphi_{\sharp}: \text{Aut}_{\mathcal{F}}(P) \rightarrow \text{Aut}_{\mathcal{F}}(P)$ , defined as  $\varphi_{\sharp}(\alpha) = \varphi \circ \alpha \circ (\varphi|_P)^{-1}$ , is fixed-point-free for all  $\varphi$ -invariant subgroup  $P \leq S$ .

The next result shows that this definition generalizes the concept of fixed-point-free automorphism of a finite group:

**Proposition 2.11.** *Let  $G$  be a finite group and  $p$  be a prime dividing the order of  $G$ . If  $\varphi: G \rightarrow G$  is a fixed-point-free automorphism then  $\varphi$  induces a fixed-point-free automorphism of  $\mathcal{F}_G(S)$ , where  $S$  is the only  $\varphi$ -invariant Sylow  $p$ -subgroup of  $G$ .*

*Proof.* According to Theorem 2.9(a) there exists  $S$  a unique  $\varphi$ -invariant Sylow  $p$ -subgroup of  $G$ . As  $\varphi|_S$  is the restriction of a fixed-point-free group morphism, it is also fixed-point-free.

Consider now  $P \leq S$  such that  $\varphi(P) = P$ . Then,  $C_G(P)$  and  $N_G(P)$  are  $\varphi$ -invariant and  $\varphi|_{C_G(P)}$  and  $\varphi|_{N_G(P)}$  are fixed-point-free. Using Theorem 2.9(b),  $\varphi$  induces a fixed-point-free group morphism

$$\varphi_{\sharp}: \text{Aut}_{\mathcal{F}_G(S)}(P) \cong N_G(P)/C_G(P) \rightarrow N_G(P)/C_G(P) \cong \text{Aut}_{\mathcal{F}_G(S)}(P).$$

□

### 3. THOMPSON THEOREMS FOR SATURATED FUSION SYSTEMS

In this section we give a unified proof of Theorems 1.1 and 1.2. In order to do so, we introduce the concept of  $\mathcal{T}$ -automorphism:

**Definition 3.1.** Let  $(S, \mathcal{F})$  be a saturated fusion system. We say that  $\mathcal{F}$  admits  $\mathcal{T}$ -automorphisms if there exists  $\mathcal{U} \leq \text{Aut}(\mathcal{F})$  such that one of the following holds:

- $\mathcal{U} = \langle \varphi \rangle$  where  $\varphi$  is a fixed-point-free automorphism of prime order.
- For every  $\mathcal{U}$ -invariant normal subgroup  $Q \trianglelefteq S$ ,  $\text{Aut}_{\mathcal{F}}(Q)$  is a  $p$ -group.

The following lemma is a particular case for finite groups which can be proved directly.

**Lemma 3.2.** *Let  $G$  be the semidirect product  $V \rtimes H$  where  $V$  is an elementary abelian  $p$ -group and  $H$  is a group with an element of order prime to  $p$  which does not centralize  $V$ . Then, given any  $p$ -syllow subgroup  $S$  of  $G$ ,  $\mathcal{F}_S(G)$  does not admit  $\mathcal{T}$ -automorphisms that leave  $V$  invariant.*

*Proof.* Assume  $G = V \rtimes H$  is a minimal counterexample to the statement,  $S$  be a  $p$ -syllow subgroup of  $G$ , and  $\mathcal{U}$  be a group of  $\mathcal{T}$ -automorphisms of  $\mathcal{F}_S(G)$  that leaves  $V$  invariant. Since  $V$  is  $\mathcal{U}$ -invariant normal in  $G$ , it is so in  $S$ : According to the hypothesis,  $\text{Aut}_{\mathcal{F}}(V) = N_G(V)/C_G(V)$  contains an element of order prime to  $p$  (coming from  $H$ ), hence we may assume that  $\mathcal{U} = \langle \varphi \rangle$  where  $\varphi$  is a fixed-point-free automorphism of prime order  $r$ . Without loss of generality, we may assume that  $\varphi$  is an honest fixed-point-free automorphism of  $G$ .

The automorphism  $\varphi$  restricts to a fixed-point-free automorphism of  $V$ , and according to Theorem 2.9(b),  $\varphi$  induces a fixed-point-free automorphism on  $G/V \cong H$ , namely  $\tilde{\varphi}$ .

Let  $q$  be any prime dividing the order of  $H$ . By Theorem 2.9(a), there exists  $Q$ , a  $\tilde{\varphi}$ -invariant Sylow  $q$ -subgroup in  $H$ . Now, if  $Q \not\leq V$ ,  $\varphi$  induces a fixed-point-free automorphism on  $V \rtimes Q \leq G$  in contradiction to the minimality of  $G$ . Therefore  $H$  must be a  $q$ -group. Moreover, if  $H$  is not abelian then the center  $Z(Q) \leq H$  is  $\tilde{\varphi}$ -invariant and  $\varphi$  induces a fixed-point-free automorphism on  $V \rtimes Z(Q) \leq G$  in contradiction to the minimality of  $G$ . Hence  $H$  is abelian. Finally if  $H$  is not elementary abelian, then characteristic subgroup  $\Omega(H) \leq H$  of elements of order  $q$  is  $\tilde{\varphi}$ -invariant and  $\varphi$  induces a fixed-point-free automorphism on  $V \rtimes \Omega(H) \leq G$  in contradiction to the minimality of  $G$ .

Now  $r$ , the order of  $\varphi$ , is different to  $p$ : the restriction of  $\varphi$  to  $V$  gives a fixed-point-free action over  $V$ , a  $p$ -group, so  $r \neq p$ .

We can assume that  $H$  acts over  $V$  without fixed points: consider a set  $A = \{h_1, \dots, h_n\}$  of generators of  $H$ . Also  $\varphi(A)$  generates  $H$ . If  $x \in V$  and  $h \in H$ , then we have the identity  $\varphi(h(x)) = \varphi(h)(\varphi(x))$ , so if  $h(x) = x$  for all  $h \in H$ ,  $\varphi(x) = \varphi(h)(\varphi(x))$ . But  $H = \{\varphi(h)\}_{h \in H}$  and the fixed points by  $H$  is a  $\varphi$  invariant subgroup. Let  $N$  be the subgroup of  $V$  of fixed points by  $H$  and assume  $N$  is not trivial. Then we can construct the group  $(V/N) \rtimes H$  and  $\varphi$  induces a fixed-point-free action by Theorem 2.9(b), and we get a contradiction with the minimality of  $G$ .

Consider now  $L$  the semidirect product  $H \rtimes \mathbb{Z}/r$ . The centralizer of  $H$  in  $L$  is itself, and we are assuming that  $V$  is a faithful  $H$ -module, and  $p \neq r$ . Then we can apply [Gor68, Theorem 3.4.4] to deduce that  $\varphi$  cannot restrict to a fixed-point-free automorphism of  $V$ , getting a contradiction.  $\square$

By [BCG<sup>+</sup>05, 4.3], if  $\mathcal{F}$  is a saturated fusion system over a  $p$ -group  $S$  and  $P$  is an  $\mathcal{F}$ -centric fully normalized in  $\mathcal{F}$  subgroup of  $S$ , there is, up to isomorphism, a unique finite group  $L = L_P^{\mathcal{F}}$  having  $N_S(P)$  as a Sylow  $p$ -subgroup such that  $C_L(P) = Z(P)$  and  $N_{\mathcal{F}}(Q) = \mathcal{F}_{N_S(P)}(L)$ .

Recall that a saturated fusion system  $\mathcal{F}$  over a  $p$ -group  $S$  is called  $H$ -free, where  $H$  is a finite group, if  $H$  is not involved in any of the groups  $L_P^{\mathcal{F}}$ , with  $Q$  running over the set of  $\mathcal{F}$ -centric fully normalized in  $\mathcal{F}$  subgroups of  $S$ .

*Proof of Theorems 1.1 and 1.2.* We will proceed considering a minimal counterexample and getting a contradiction. So, let  $S$  be the smallest  $p$ -group and  $\mathcal{F}$  the saturated fusion system with a minimal number of morphisms such that  $(S, \mathcal{F})$  admits  $\mathcal{U}$  a group of  $\mathcal{T}$ -automorphisms and  $\mathcal{F}_S(S) \not\leq \mathcal{F}$ .

**Step 1:** There exists a non-trivial elementary abelian proper subgroup  $W(S) \leq S$  such that  $(S, \mathcal{F}) = (S, N_{\mathcal{F}}(W(S)))$ :

Given a group  $G$ , let  $Z(G)$ ,  $J(G)$ , and  $\Omega(G)$  denote the center, the Thompson subgroup, and the group generated by the elements of order  $p$  of  $G$  respectively. Let  $W(S)$  be the characteristic subgroup of  $S$  defined in [OS09, Section 4]. Then  $\Omega(Z(S)) \leq W(S) \leq \Omega(Z(J(S)))$ , and  $W(S)$  is non-trivial and elementary abelian. Moreover, as  $W(S)$  is characteristic, it is  $\mathcal{U}$ -invariant and normal in  $S$ . This implies that  $W(S)$  is fully normalized in  $\mathcal{F}$  so, by Proposition 2.3,  $N_{\mathcal{F}}(W(S))$  is a saturated fusion system over  $S$ .

Let us see now that  $(S, \mathcal{F}) = (S, N_{\mathcal{F}}(W(S)))$ : assume  $N_{\mathcal{F}}(W(S)) \not\leq \mathcal{F}$ . Then  $\mathcal{U}$  induces a group of  $\mathcal{T}$ -automorphisms in  $(S, N_{\mathcal{F}}(W(S)))$ , so if there are morphisms in  $\mathcal{F}$  which are not in  $N_{\mathcal{F}}(W(S))$ , by minimality of  $\mathcal{F}$ ,  $N_{\mathcal{F}}(W(S)) = \mathcal{F}_S(S)$ . But, as  $p$  is an odd prime (respectively  $p = 2$  and  $\mathcal{F}$   $\Sigma_4$ -free), by [OS09, Theorem 1.3] (respectively [OS09, Theorem 1.1]) this implies  $\mathcal{F} = \mathcal{F}_S(S)$ , so  $N_{\mathcal{F}}(W(S)) = \mathcal{F}$ , getting a contradiction.

If  $W(S) = S$  then, applying Lemma 3.2,  $(S, \mathcal{F}) = (S, \mathcal{F}_S(S))$ , so it is not a counterexample.

**Step 2:** According to Lemma 2.4,  $(S, \mathcal{F})$  projects onto a saturated fusion system  $(S/W(S), \mathcal{F}/W(S))$  and  $\mathcal{U}$  induces a group of  $\mathcal{T}$ -automorphism of  $\mathcal{F}/W(S)$ . So, by the minimality hypothesis,  $\mathcal{F}/W(S) = \mathcal{F}_{S/W(S)}(S/W(S))$ .

**Step 3:** There is an element  $\alpha$  of prime order  $q$  in  $\text{Aut}_{\mathcal{F}}(W(S))$ , with  $q \neq p$ :

As we are assuming that  $\mathcal{F}_S(S) \not\leq \mathcal{F}$ , using the Alperin's Theorem for saturated fusion systems (Theorem 2.6), there exists  $P$ , a fully normalized in  $\mathcal{F}$ ,  $\mathcal{F}$ -centric,  $\mathcal{F}$ -radical subgroup of  $S$  with an  $\mathcal{F}$ -automorphism  $\bar{\alpha}$  of prime order  $q$ ,  $q \neq p$ . By [BCG<sup>+</sup>05, Proposition 1.6], as  $W(S)$  is normal in  $\mathcal{F}$  and  $P$  is  $\mathcal{F}$ -radical,  $W(S)$  is contained in  $P$  and we have induced maps in the normal series  $1 \trianglelefteq W(S) \trianglelefteq P$ . By Step 2  $\bar{\alpha}$  projects to the identity in  $P/W(S)$ . If the restriction of  $\bar{\alpha}$  to  $W(S)$  is also the identity, by [Gor68, Theorem 5.3.2] the order of  $\bar{\alpha}$  is a power of  $p$ , getting a contradiction. So  $\bar{\alpha}$  restricts to an automorphism  $\alpha$  on  $W(S)$  of order  $q$ .

**Last step:** We finish the proof by considering  $G = W(S) \rtimes H$  where  $H = \text{Aut}_{\mathcal{F}}(W(S))$ , then the automorphism group induced by  $\mathcal{U}$  on  $G$  is a group of  $\mathcal{T}$ -automorphism that leaves  $W(S)$  invariant, what contradicts Lemma 3.2. Therefore there is no minimal counterexample  $(S, \mathcal{F})$ .  $\square$

We finish this section with a couple of examples that illustrate the scope of these results. The first example, what seems to be well known for the experts (see comments below [Tho59, Theorem A]), shows that the  $\Sigma_4$ -free hypothesis in the  $p = 2$  case is a necessary one.

**Example 3.3.** Let  $G$  be the simple group  $L(2, 17)$ , and let  $S \leq G$  be a 2-sylow subgroup. According to [CCN<sup>+</sup>85, p. 9], the 2-fusion is completely determined by its self normalizing sylow  $S \cong D_{16}$  and two conjugacy classes of rank two elementary abelian subgroups of type  $2A^2$  whose normalizer is isomorphic to  $\Sigma_4$ . So  $\mathcal{F} = \mathcal{F}_S(G)$  is not  $\Sigma_4$ -free,  $\mathcal{F} \neq \mathcal{F}_S(S)$ , and for every normal  $Q \trianglelefteq S$ ,  $\text{Aut}_{\mathcal{F}}(Q)$  is a 2-group (since the only maximal subgroup of  $G$  of index coprime to 2 is  $S$ ). Therefore, for every  $\mathcal{U} \leq \text{Aut}(\mathcal{F})$ , and  $\mathcal{U}$ -invariant normal subgroup  $Q \trianglelefteq S$ ,  $\text{Aut}_{\mathcal{F}}(Q)$  is a 2-group, but  $\mathcal{F} \neq \mathcal{F}_S(S)$ .

The last example shows that group automorphisms with fixed points can induce a fixed-point-free automorphisms of saturated fusion system.

**Example 3.4.** Let  $S$  be a finite  $p$ -group such that there exist a fixed-point-free automorphism  $\phi$  of  $S$ . Consider  $H = S \rtimes \langle \phi \rangle$  and let  $N$  be a group whose order is coprime to  $p$  and a homomorphism  $f: H \rightarrow \text{Aut}(N)$ . The homomorphism  $f$  defines a semidirect product  $G = N \rtimes H$ , where the subgroup  $K = SN$  is a normal subgroup of  $G$  and  $\phi$  acts on  $K$  not necessary without fix points, but it acts on the fusion category  $\mathcal{F}_S(G) = \mathcal{F}_S(S)$  without fix points.

An example of this situation is the group  $A_4$  and  $\phi \in \text{aut}(A_4)$  given by conjugation in  $S_4$  by the transposition  $(1, 2)$ . Then  $\phi$  fixes the Sylow 3-subgroup  $S = \langle (1, 2, 3) \rangle$ , and furthermore it acts without fix points on the fusion category  $\mathcal{F}_S(A_4)$ .

## REFERENCES

- [BCG<sup>+</sup>05] Carles Broto, Natàlia Castellana, Jesper Grodal, Ran Levi, and Bob Oliver. Subgroup families controlling  $p$ -local finite groups. *Proc. London Math. Soc.* (3), 91(2):325–354, 2005.
- [BLO03] Carles Broto, Ran Levi, and Bob Oliver. The homotopy theory of fusion systems. *J. Amer. Math. Soc.*, 16(4):779–856, 2003.

- [CCN<sup>+</sup>85] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson. *Atlas of finite groups*. Oxford University Press, Eynsham, 1985. Maximal subgroups and ordinary characters for simple groups, With computational assistance from J. G. Thackray.
- [CSV13] Jose Cantarero, Jerome Scherer, and Antonio Viruel. Nilpotent  $p$ -local finite groups. *Ark. Mat.*, 2013.
- [DGPS11] Antonio Díaz, Adam Glesser, Sejong Park, and Radu Stancu. Tate’s and Yoshida’s theorems on control of transfer for fusion systems. *J. Lond. Math. Soc. (2)*, 84(2):475–494, 2011.
- [Gor68] Daniel Gorenstein. *Finite groups*. Harper & Row Publishers, New York, 1968.
- [KL08] Radha Kessar and Markus Linckelmann.  $ZJ$ -theorems for fusion systems. *Trans. Amer. Math. Soc.*, 360(6):3093–3106, 2008.
- [Oli04] Bob Oliver. Equivalences of classifying spaces completed at odd primes. *Math. Proc. Cambridge Philos. Soc.*, 137(2):321–347, 2004.
- [OS09] Silvia Onofrei and Radu Stancu. A characteristic subgroup for fusion systems. *J. Algebra*, 322(5):1705–1718, 2009.
- [RV04] Albert Ruiz and Antonio Viruel. The classification of  $p$ -local finite groups over the extraspecial group of order  $p^3$  and exponent  $p$ . *Math. Z.*, 248(1):45–65, 2004.
- [Tho59] John Thompson. Finite groups with fixed-point-free automorphisms of prime order. *Proc. Nat. Acad. Sci. U.S.A.*, 45:578–581, 1959.
- [Tho60] John G. Thompson. Normal  $p$ -complements for finite groups. *Math. Z.*, 72:332–354, 1959/1960.
- [Tho64] John G. Thompson. Normal  $p$ -complements for finite groups. *J. Algebra*, 1:43–46, 1964.

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE PAÍS VASCO, APARTADO 644, 48080 BILBAO, SPAIN.

*E-mail address:* jon.gonzalez@ehu.es

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 Cerdanyola del Vallès, SPAIN.

*E-mail address:* Albert.Ruiz@uab.cat

DEPARTAMENTO DE ÁLGEBRA, GEOMETRÍA Y TOPOLOGÍA, UNIVERSIDAD DE MÁLAGA, CAMPUS DE TEATINOS, S/N, 29071-MÁLAGA, SPAIN.

*E-mail address:* viruel@agt.cie.uma.es